




# Experience Flow




### Sales/Marketing - Preordering






#### Collection




• The instruments and assays used by our partner laboratories are comparable to those used in

#### Results


everlywell Shop Tests ~ The Science Help ~ My Everly Welcome, Rebecca	Register Your Kit <u>My Everly</u> Account ✓	everlywell       Shop Tests 、       The Science       Help 、       Register Your Kit       My Everly       Account 、         < Dashboard       Get \$35!       Save a PDF 、        Share 、	<ul> <li>Comparison of the state of the</li></ul>			7 Foods foods (Class 1 foods ion, but they can be s
<section-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header><image/><image/></section-header></section-header></section-header></section-header></section-header></section-header></section-header></section-header>		<image/> <image/> <section-header><section-header><section-header><section-header><section-header><section-header><section-header><text><text><text><text></text></text></text></text></section-header></section-header></section-header></section-header></section-header></section-header></section-header>	diet. Moderate Reactivity: 3 Foods Moderate reactivity foods (Class 2 foods in symptoms as well. Coffee Cow's Milk Yogurt Normal Reactivity T About coffee Coffee contains caffeir measuring your body's content. If reactive to a elimination diet. A good Hidden Sources Coffee bean chocolate	ar other reactivity levels if you decide to do an elimination In your lab results) can cause Toderate Reactivity Toderate Reactivity Mild Reactivity 50 Reactivity 16 High Reactivity the, which is a naturally occurring stimulant. This test is a IgG antibody response to the coffee bean, not the caffeine coffee, it's recommended to be avoided during a temporary ad substitute could be matcha or other herbal teas. es, decaf coffee, certain desserts such as tiramisu, cakes.	All foods will o were measure but if you're co	Your Eggplant Normal Reactivity About eggplan Eggplant is a ve can be prepare (eggplant dip). during a tempo Hidden Source Ratatouille, Bak as stir fries and as stir fries and cause some amount of ed in the normal rang urious, you can see the nal Reactivity Foods
Results Landing Page		Test Specific Page	Results from High to low			




**Related Resources and Information** 



## Sample Reports





88.5	44.3
30.6	15.5
0	c
Evening	Nigh
~ -	
	Č Ever

Cortisol >	DHEAS 6.7	>	Estradiol	2.0	FSH	6.1
		na				••••••••••••••••••••••••••••••••••••••
	6.7 mcg/dL		2 pg/mL		6.1 U/L	Normal 2,4+1
Free	ГЗ	> Free	T4	> LH		>
		4.0	1.6		5.9	
	ii.			23		8.7
4 pg/n		iormai 2.4 - 4.2 1.6 n		rmai 0.7 - 2.5 5,9 L		irmai 1.6-9.3
Proge	sterone	> Prog	gesterone/Estradio	I Ratio > TPO	ab	×
	166.0		3.0		13.0	
-	84	275.0	1001	500.3		710
166 pj	g/mL Nom	nal 75,0~270,0 83		Low < 100.0 13 II.	l/mL )	Norma) <70.0
	TS		> Tes	tosterone	>	
		1,9		35.0		
	-	04	34		lan	
		) µLi/mi. Nor			al 16.0 - 55.0	

